

National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Improving First-Guess Surface IR Emissivity Models

AIRS Science Team Meeting
May 3-6, 2005

Evan Fishbein
California Institute of Technology
Jet Propulsion Laboratory

5 May 2005




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Why Improve Emissivity First-Guess?

- **Currently large component of final solution is from first guess**
- **Cloud-clearing maps inter-footprint surface variability into cloud field**
 - Include footprint-dependent surface in Obs-Calc for CC
- **Reduce degree-of-freedom by coupling emissivity at different frequencies**
- **Improve convergence rate**




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Current Surface Emissivity First Guess

- **Final retrieval initialized with surface emissivity regression**
- **Surface emissivity training set from AIRS Level 2 Simulation System (AL2SS)**
- **AL2SS surface model was not designed for this purpose**
 - Provides worse-case surface variability
 - Statistics are not representative
 - Uniform skin temperature over footprint




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



AL2SS Surface Model

- Contains
 - Two “soil” types: quartz sand and urban sprawl
 - Three vegetation types: conifer, deciduous, grass
 - Constant salinity sea water
 - Two types of ice: ice and snow
- Each footprint is a mixture of components based on
 - Interpolated 1km NDVI at center of footprint
 - (not averaged over AIRS footprint)
 - 1 km simplified IGBP Global Land-Cover Type
 - Vegetation type randomly distributed among possible vegetation types on
- Lambertian emissivity (except sea water)
- Constant skin temperature




National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Material Emissivity Models

- Extrapolated shortwave emissivities
- Quartz has anomalous Reistrahlen band
- Ice emissivity depends on grain size and water content






National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California




Global Land Cover Characterization

International Geosphere Biosphere Programme Global Land Cover Type




1. Evergr. Needle For.	7. Open Shrub.	13. Urban
2. Evergr. Broad For.	8. Woody Savannas	14. Crop/Mosaic
3. Decid. Needle For.	9. Savannas	15. Snow/Ice
4. Decid. Broad For.	10. Grassland	16. Barren/Desert
5. Mixed Forest	11. Wetlands	17. Water
6. Closed Shrub.	12. Crops	18. Tundra



- Actively developed by MODIS Land Science Team
- Types of vegetation relatively unimportant

AIRS Science Team Meeting: 05-May-05-6



National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California




Soil Variability

Soil Classification

- Organic material
- Weathering



AIRS Science Team Meeting: 05-May-05 - 7




National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Rock Variability



AIRS Science Team Meeting: 05-May-05 - 8





National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Vegetation





National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Observed Variability

- **ASTER**


Swath width: 60km

Spatial Resolution: 90m


- **MODIS**

Swath width: 2330 km


Spatial Resolution: 1000 m




AIRS Science Team Meeting: 05-May-05 - 10



ASTER Emissivity






National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



AIRS ASTER Intercomparison

8.65 μm






National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Factors Affecting First Guess Emissivity

- Amount of vegetation and view of soil
- Soil composition and particle size
- Surface type - snow / ice, vegetation type
- Soil moisture



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



First Guess Options

1. Construct spectral emissivity map

- Monthly L3 product (possibly higher spatial resolution)
- Emissivity maps from ASTER/MODIS
- Issues
 - Ignore non-Lambertian behavior
 - Spectral coverage
 - Vegetation annual and seasonal variability

2. Evaluate emissivity using extended simulation framework

- More components and means of estimating components
- Spectral model of components
- Issues
 - Soil composition model
 - Vegetation in polar winter
 - Can components be derived from long wave emissivity
 - Use MODIS/ASTER emissivities to determine composition

AIRS Science Team Meeting: 05-May-05 - 14



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California




Using Land Cover Type to Organize Activity

Easy

- **Tropical Deciduous forests**
 - High vegetation cover
 - Weak seasonal variability
 - Lambertian
- **Conifer Forest**
 - Moderate vegetation cover
 - Weak seasonal dependence
 - Some non-Lambertian behavior
- **Mid-latitude Deciduous forests**
 - High vegetation cover
 - Strong seasonal dependence
 - Lambertian (?)

Difficult

AIRS Science Team Meeting: 05-May-05 - 15



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Using Land Cover Type to Organize Activity

Hard



- **Grasslands**

- Low vegetation cover
- Strong seasonal variability
- Non-Lambertian

- **Marsh**


- Variable vegetation
- Strong seasonal variability (vegetation and soil)
- Lambertian (?)

- **Desert**

- No vegetation
- Very weak seasonal variability
- Lambertian (?)

Harder

AIRS Science Team Meeting: 05-May-05 - 16



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Using Land Cover Type to Organize Activity

Hard



- **Tundra**

- Low vegetation cover
- Diurnal, weekly, seasonal variability
- Snow, ice and frozen soil
- Lambertian (?)

- **Snow / Ice**

- Diurnal, weekly, seasonal variability
- Non-Lambertian (?)
- Microwave even more complicated

- **Urban (?)**


- **Varied Topography**

- Variable surface pressure and water vapor
- Heterogeneous soil composition



**Good
Luck**

AIRS Science Team Meeting: 05-May-05 - 17



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



References

- Jacob, F, Petitcolin, F, Schmugge, T, Vermote, E, French, A. and Ogawa, K, *Comparison of land surface emissivity and radiometric temperature derived from MODIS and ASTER sensors*, **Rem. Sens. Environ**, 90, 137–152, 2004.
- Peres, L.F. and DaCamara, C.C, *Land surface temperature and emissivity estimation based on the two-temperature method: sensitivity analysis using simulated MSG/SEVIRI data*, **Rem. Sens. Environ**, 91, 377-389, 2004.
- Snyder, W.C, Wan, Z, Zhang, Y. and Feng, Y-Z, *Classification-based emissivity for land surface temperature measurement from space*, **Int. J. Rem. Sens**, 19, 2753-2774, 1998.
- Szciech-Gajewska, M. and Rabier, M, *Surface emissivity for use in assimilation of IR radiance data over land*, **Proc. Intl TOVS Study Conf**, 13, 241-249, 2003.
- Wan, Z-M and Li, Z-L, *Physics-based algorithm for retrieving land-surface emissivity and temperature from EOS/MODIS Data*, **IEEE Trans. Geosci**, 35, 980-996. 1997.
- Wan, Z, Zhang, Y, Zhang, Q. and Li, Z-l, *Validation of the land-surface temperature products retrieved from Terra Moderate Resolution Imaging Spectroradiometer data*, **Rem. Sens. Environ**, 83, 163–180, 2002.